

Chern-Simons Gravity, Wilson Lines and Large N Dual Gauge Theories

L.D. Paniak

Joseph Henry Laboratories, Department Of Physics

Princeton University

Princeton, New Jersey 08544, USA

e-mail: paniak@feynman.princeton.edu

Abstract

A five-dimensional Chern-Simons gravity theory based on the anti-de Sitter group $SO(4, 2)$ is argued to be a useful model in which to understand the details of holography and the relationship between generally covariant and dual local quantum field theories. Defined on a manifold with boundary, conformal geometry arises naturally as a gauge invariance preserving boundary condition. By matching thermodynamic quantities for a particular background geometry, the dimensionless coupling constant of the Chern-Simons theory is directly related to the number of fields in a putative dual theory at high temperature. As a consistency check, the semiclassical factorization of Wilson line observables in the gravity theory is shown to induce a factorization in dual theory observables as expected by general arguments of large N gauge theory.

Holography in D -dimensional generally covariant system is the general proposition which states that the number of degrees of freedom in the system is bounded from above by the area (in Planck units) of a $D - 2$ -dimensional spatial surface surrounding the system [1, 2, 3]. An ambitious extension of this concept is the organization of the so-called boundary degrees of freedom into a local quantum field theory which effectively describes the physics of the D -dimensional system. The existence of such ‘dual’ theories to certain generally covariant systems has been supported by much recent work, especially in the case of compactified string theory and its relation to supersymmetric gauge theory. Initial evidence from D-brane scattering calculations [4, 5] has been formalized in the ‘AdS/CFT’ correspondence principle [6, 7, 8] (see [9] for a comprehensive review with exhaustive reference list). Unfortunately, due to the complexities of explicit calculations on both the string and gauge theory sides of the duality, the details of the relationship between these theories remain uncertain.

Since holography is considered to be a generic feature of generally covariant systems and it can be argued theories which admit anti-de Sitter space-times also admit dual interpretations [10], it may be useful to examine examples outside of string theory with these properties. With these conditions in mind, we propose to examine a generally covariant system in five space-time dimensions which is based on a pure Chern-Simons gauge theory. With the particular choice of the anti-de Sitter group $SO(4, 2)$ as the gauge group, the Chern-Simons theory defines a higher curvature generalization of Einstein gravity [11, 12, 13] which naturally admits anti-de Sitter space-times. In addition, when defined on space-times with a boundary, the natural gauge invariance preserving boundary condition of a flat connection immediately leads to a boundary manifold on which unrestricted gauge transformations generate conformal symmetries. In this way, the Chern-Simons gravity theory is similar to low-energy string theory effective actions which admit anti-de Sitter backgrounds with conformally invariant boundaries, but there are differences. Unlike higher curvature terms which are expected due to stringy α' corrections to Einstein gravity, in Chern-Simons gravity higher curvatures are not suppressed relative to the linear curvature term. Consequently, the Chern-Simons gravity theory we will discuss here does not follow in an obvious way as the low energy effective description of a string theory configuration. Unfortunately this means we will not have at our disposal the well-known machinery to identify the conjectured dual gauge theory. Moreover, since Chern-Simons gravity is not renormalizable in five space-time dimensions we will restrict ourselves

to the semiclassical regime and consider the theory in the vicinity of solutions of the classical equations of motion.

While it may seem paradoxical to use a gauge theory to define a gravitational theory, the Chern-Simons gravity theory we will consider is a direct generalization of the $2 + 1$ -dimensional theory which describes Einstein gravity in three dimensions [14, 15]. As in the three-dimensional case, the relationship to a gauge theory leads to the most intriguing property of a Chern-Simons formulation of gravity - Wilson line observables. These are natural gauge-invariant observables in the Chern-Simons theory and are carried over to the gravity theory as a concrete and effective definition of non-local, diffeomorphism-invariant observables. Assuming there exists a dual (holographic) theory which describes the same physics as the gravitational theory, such observables should have a natural interpretation in terms of observables of a local quantum field theory.

$$\langle X \rangle_{CS} = \langle Y \rangle_{QFT} \quad (1)$$

Here we will show explicitly this relationship implies the factorization of field theory observables consistent with the expectations for large rank gauge theories. This will be accomplished in the semiclassical regime of the Chern-Simons gravity theory by finding a specific background geometry which is argued to control the system at high temperatures. The matching of free energies between the gravitational theory and a hot, rank N gauge theory leads to an identification between the number of gauge degrees of freedom and the coupling constant of the Chern-Simons gravity theory. In this way, the large N limit of the gauge theory is shown to correspond to the semiclassical limit of the gravity theory and the factorization of observables follows.

Chern-Simons gravity is a higher curvature generalization of Einstein gravity [11, 12, 13] based on a five-dimensional pure Chern-Simons theory with anti-de Sitter gauge group, $SO(4, 2)$. The five-dimensional Chern-Simons form is defined via the relation

$$d\omega_{CS} = \langle F \wedge F \wedge F \rangle \quad (2)$$

Here F is the curvature two-form related to the gauge connection A in the familiar way

$$F = dA + A \wedge A \quad (3)$$

The bracket $\langle \cdots \rangle$ defines an invariant trace operation on the gauge algebra. In general, it is not unique and, in particular, depends on the representation of the gauge group chosen. The choice which will be used in the sequel is that of the fully anti-symmetric tensor invariant of $SO(4, 2)$

$$\langle J_{B_1 C_1} J_{B_2 C_2} \cdots J_{B_n C_n} \rangle = \epsilon_{B_1 C_1 B_2 C_2 \cdots B_n C_n} \quad (4)$$

Here J_{BC} are generators of the gauge group and $B, C = 1, \dots, 6$ label $so(4, 2)$ algebra indices. The Chern-Simons form can be explicitly calculated in terms of the connection

$$\omega_{CS} = \langle A \wedge dA \wedge dA + \frac{3}{2} A \wedge A \wedge A \wedge dA + \frac{3}{5} A \wedge A \wedge A \wedge A \wedge A \rangle \quad (5)$$

The action is simply the integral of this five-form over a given manifold $\mathcal{M}$

$$S = k \int_{\mathcal{M}} \omega_{CS} \quad (6)$$

where k is a dimensionless coupling constant. We will restrict ourselves to topologically trivial $\mathcal{M}$ for simplicity.

Of primary interest are manifolds with boundary. In order to maintain gauge invariance of the action on a manifold with boundary, it is necessary to impose boundary conditions on the gauge fields and gauge transformations. Under a gauge transformation

$$A_g = g^{-1} A g + g^{-1} dg \quad (7)$$

the Chern-Simons form transforms as

$$\omega_{CS}(A_g) = \omega_{CS}(A) + d\alpha(A, g) + \frac{1}{10} \langle (g^{-1} dg)^5 \rangle \quad (8)$$

The last term in (8) has the form of a winding number and upon integration over $\mathcal{M}$ takes values in the fifth cohomology group $H^5(\mathcal{M}, \pi_5(SO(4, 2)))$. Since $\pi_5(SO(4, 2))$ is pure torsion the cohomology group is trivial and there is no quantization of the coupling constant k . Consequently, for manifolds without boundary the action is gauge invariant. With a boundary, there generically is a non-vanishing contribution from the four-form α

$$\alpha(A, g) = \langle -\frac{1}{2} dg g^{-1} (AF + FA) + \frac{1}{2} dg g^{-1} A^3 + \frac{1}{4} (dg g^{-1} A)^2 + \frac{1}{2} (dg g^{-1})^3 A \rangle \quad (9)$$

Gauge invariance requires the integral of α over the four- dimensional boundary $\partial\mathcal{M}$ to vanish. There are a number of ways to achieve this result. One way is to follow as in the

three-dimensional case [17] and restrict the gauge transformations allowed at the boundary to be trivial and leave the gauge field unconstrained. The restriction we will use here is different and has an appealing interpretation in terms of a conformal geometry induced on the boundary. We will constrain the curvature F to vanish on the boundary which allows one to choose the connection A to be pure gauge. Gauge transformations at the boundary are left unrestricted. It can be shown that under this restriction the Chern-Simons form transforms invariantly up to winding terms which we have previously argued are irrelevant for the gauge group $SO(4, 2)$.

Having specified the action and associated boundary conditions on the fields, the equations of motion follow from a bulk variation of the action (6) and are conveniently given by

$$\epsilon_{B_1 C_1 B_2 C_2 B_3 C_3} F^{B_1 C_1} \wedge F^{B_2 C_2} = 0 \quad (10)$$

On a manifold with boundary, the variational principle for an action is typically ill-defined due to boundary contributions. In the present case the boundary contribution can be calculated as

$$\int_{\partial \mathcal{M}} \langle A \wedge A \wedge A \wedge \delta A \rangle \quad (11)$$

Fortunately, the choice of flat boundary conditions can be argued to eliminate these contributions [16].

The relation of the Chern-Simons action to a theory of gravitation follows from decomposing the connection A on a basis of the anti-de Sitter algebra $\{P_a, J_{ab}\}$.

$$A = \frac{e^a}{\lambda} P_a + \omega^{ab} J_{ab} \quad (12)$$

The coefficients of this expansion are identified with the fünfbein and spin connection

$$A^{ab} = \omega^{ab} \quad , \quad A^{a5} = \frac{e^a}{\lambda} \quad a, b = 0, 1, \dots, 4 \quad (13)$$

An extra length scale parameter λ , related to the cosmological constant, has been introduced in the definition of the fünfbein. These definitions lead to the following identifications for the gauge field curvature in terms of geometric curvature and torsion

$$\begin{aligned} F^{ab} &= d\omega^{ab} + \omega^{ac} \wedge \omega_c^b + \frac{e^a \wedge e^b}{\lambda^2} \equiv R^{ab} + \frac{e^a \wedge e^b}{\lambda^2} \\ F^{a5} &= de^a + \omega_c^a \wedge e^c \equiv T^a \end{aligned} \quad (14)$$

The flatness condition $F = 0$ from the gauge theory translates to a restriction to torsion-free configurations with a specific curvature two-form

$$R^{ab} + \frac{e^a \wedge e^b}{\lambda^2} = 0 \quad , \quad T^a = 0 \quad (15)$$

This curvature two-form describes a space with constant, negative curvature - an anti-de Sitter space - and we see our choice of gauge invariance preserving boundary conditions $F = 0$ produces the natural setting for the duality between a bulk gravitational theory and some local quantum field theory associated with the boundary. As is well-known by now, taking the boundary of the manifold to be defined as the surface on which the anti-de Sitter metric becomes singular induces a conformal structure on the boundary $\partial\mathcal{M}$. Moreover, the unrestricted $SO(4, 2)$ gauge freedom at the boundary acts on this space as conformal motions [8].

The result of substituting the decomposition of the gauge field (14) into the Chern-Simons form (5) is a higher curvature generalization of the Einstein-Hilbert action which has been investigated previously [11, 12, 13]

$$S = k\epsilon_{abcde} \int_{\mathcal{M}} \left[\frac{e^a \wedge R^{bc} \wedge R^{de}}{\lambda} + \frac{2e^a \wedge e^b \wedge e^c \wedge R^{de}}{3\lambda^3} + \frac{e^a \wedge e^b \wedge e^c \wedge e^d \wedge e^e}{5\lambda^5} \right] \quad (16)$$

and the equations of motion in gravitational variables are given by

$$\begin{aligned} \epsilon_{abcde} \left(R^{ab} + \frac{e^a \wedge e^b}{\lambda^2} \right) \wedge \left(R^{cd} + \frac{e^c \wedge e^d}{\lambda^2} \right) &= 0 \\ \epsilon_{abcde} \left(R^{ab} + \frac{e^a \wedge e^b}{\lambda^2} \right) \wedge T^c &= 0 \end{aligned} \quad (17)$$

Since Chern-Simons theory in five dimensions is not renormalizable, we restrict ourselves to the semiclassical approximation where the coupling constant k is large. Hence, we will be interested in the behaviour of the theory in the vicinity of solutions of the classical equations of motion. Immediately the symmetry of the theory suggests a background which is consistent with the boundary condition, namely anti-de Sitter space. Unlike the more familiar cases of low energy approximations to string theory, here the anti-de Sitter background is not physically appealing. To see that the Chern-Simons gravity theory is pathological in this background one needs to look no further than the classical equations of motion (10) and recall anti-de Sitter space corresponds to vanishing curvature in the gauge theory. Since the equations of motion are quadratic in the curvature F , all fluctuations about the anti-de Sitter

background vanish identically. In terms of an effective quantum theory about this background it means there are no propagating degrees of freedom and the theory is perturbatively sterile. Moreover, when calculating observables of Chern-Simons gravity we would find they are all trivial (up to topology of the space-time). In short, Chern-Simons gravity in the anti-de Sitter background cannot describe any non-trivial field theory.

Fortunately, other solutions of Chern-Simons gravity are known to exist. Of most interest to the current situation are those which are analogous to asymptotically anti-de Sitter Schwarzschild black holes in Einstein gravity [18]. These have been discussed in detail for Chern-Simons gravity in [19] and for similar theories in [20]. In particular, we are interested in a black hole, with mass M related to a dimensionless parameter a , defined by the line-element

$$ds^2 = -(1 - a + (r/\lambda)^2)dt^2 + \frac{dr^2}{1 - a + (r/\lambda)^2} + r^2 d\Omega^2 \quad (18)$$

where $d\Omega^2$ is the volume element of the three-sphere. It is easily checked that this solution of (17) leads to a flat Chern-Simons connection on the boundary at infinity radius in accordance with our boundary conditions. Since this solution represents a black hole geometry, there is a natural temperature associated with the system which can be calculated by continuing to Euclidean time with the result [19, 21]

$$\beta = \frac{1}{T} = \frac{2\pi\lambda}{\sqrt{a-1}} \quad (19)$$

As well, the Euclidean action of Chern-Simons gravity can be calculated in this background relative to zero temperature and vanishing mass $M(a)$ at $a = 1$

$$I_E = \beta M(a) - \frac{16\pi^2\lambda}{3\beta} \left(3 + \frac{4\pi^2\lambda^2}{\beta^2}\right) \quad (20)$$

Important to note is that as the temperature is increased, the contribution of thermal gravitons [22] to the action in the second term of (20) becomes arbitrarily negative. Therefore, it seems reasonable to conclude for sufficiently high temperature that the background (18) will dominate the semiclassical calculation of the partition function. This is completely analogous to the domination of the partition function at high temperatures by the Schwarzschild anti-de Sitter black hole as investigated in quantum gravity by Hawking and Page [18] and Witten [8, 23] in the context of the AdS/CFT correspondence. Consequently, at high temperatures the partition function and free energy of the system can be easily calculated semiclassically

with the result

$$\beta F_{CS} = -\log Z = k I_E \quad (21)$$

If there exists a dual, local quantum field theory defined on the boundary of the five-dimensional space-time which is described by Chern-Simons gravity in the background (18) then we should compare the free energy to that of a high temperature quantum theory. In general we expect the boundary theory to contain gauge fields and in the limit of high temperature we expect the system is in a (quark-gluon) plasma phase where fundamental degrees of freedom are not confined. If the gauge group is of rank N there are order N^2 gluonic degrees of freedom and possibly as many matter degrees of freedom. An estimate of the free energy of such a system on a four-sphere of length scale λ can be given by dimensional arguments

$$F_{QFT} = -\sigma N^2 \lambda^3 T^4 \quad (22)$$

where σ is a proportionality constant of order one. Equating the free energies in (21) and (22), we find in the limit of high temperature the Chern-Simons coupling constant is directly related to the number of degrees of freedom in the boundary field theory

$$k = \frac{3\sigma}{64\pi^4} N^2 \quad (23)$$

This identification of the coupling constant with the number of degrees of freedom is reminiscent of the work of [24] and [25] on eleven-dimensional Chern-Simons gravity and M-theory. In that case the degrees of freedom were taken to lie in the gravitational theory, not its dual and the relation to the gravitational coupling constant followed from Mach's Principle. Like that case, our arguments only hold if the number of degrees of freedom N is large: Mach's Principle requires a uniform background and we have done all our calculations in the semiclassical limit of large k which, by (23), is equivalent to large N .

A novel consistency check of the relationship between Chern-Simons gravity and an unknown, dual, local quantum field theory which describes the same physics can be formulated in terms of the observables of each theory. Using the relationship (23), we will derive the large N factorization of gauge-invariant observables in the dual theory from semiclassical gravity calculations.

While it is not understood in general how to define observables in a generally covariant theory, Chern-Simons gravity contains a natural family of observables in Wilson lines inherited

from its gauge theory formulation

$$W_R[\gamma] = \text{Tr}_R P e^{i \oint_\gamma A_\mu dx^\mu} = \text{Tr}_R P e^{i \oint_\gamma (e_\mu^a P_a / \lambda + \omega_\mu^{ab} J_{ab}) dx^\mu} \quad (24)$$

For finite-dimensional representations R of $SO(4, 2)$, this expression is simply the trace of the holonomy around the closed path γ . Of course for a non-compact gauge group like $SO(4, 2)$ there are infinite dimensional irreducible representations as well. In this case, the naive picture breaks down and the trace of the path ordered matrix integral must be realized as a dynamical system of infinite degrees of freedom [26, 27]. Physically, these observables describe the trajectories of external test particles in the gravitational background. As in a compact gauge theory, the invariant properties of these test particles are given in terms of the irreducible representations of the gauge group which for $SO(4, 2)$ are labelled by mass and spin.

The expectation value of a collection of Wilson lines is given by the standard formula

$$\langle W_{R_1}[\gamma_1] \cdots W_{R_n}[\gamma_n] \rangle_{CS} = \frac{1}{Z} \int \mathcal{D}A e^{ik \int_{\mathcal{M}} \omega_{CS}[A]} \text{Tr}_{R_1} P e^{i \oint_{\gamma_1} A_\mu dx^\mu} \cdots \text{Tr}_{R_n} P e^{i \oint_{\gamma_n} A_\mu dx^\mu} \quad (25)$$

If one assumes there exists a dual local quantum field theory to the Chern-Simons gravity, this expectation value has an alternate interpretation in terms of some gauge invariant quantity, V defined in the dual theory

$$\langle W_{R_1}[\gamma_1] \cdots W_{R_n}[\gamma_n] \rangle_{CS} = \langle V_{R_1, \dots, R_n}[\gamma_1, \dots, \gamma_n] \rangle_{QFT} \quad (26)$$

Unfortunately, there are at least two difficulties with this identification in the present situation. First, the Wilson lines in the Chern-Simons gravity theory, being observables in a generally covariant theory cannot depend on the geometry of paths $\{\gamma_i\}$. They should only be sensitive to the topological properties of the paths. This is not what one expects from a dual local quantum gauge theory in which similar path-dependent objects would be generic. In those cases one expects the Wilson line would depend on the length, or enclosed area of the paths. Second, the calculation of such observables in the Chern-Simons gravity theory appears to be beyond reach in the general case and the conjectured dual local field theory is completely unknown.

Fortunately, in the semiclassical limit of large k these concerns are absent. Leading contributions to expectation values come from the vicinity of definite geometries satisfying both

the gauge invariance preserving boundary condition, $F = 0$ and the classical equations of motion (17). In this way the topological nature of the Wilson lines is broken and we expect these observables (25) to depend on geometric information. In particular, in the limit of high temperature we have argued the black hole geometry (18) dominates the semiclassical partition function. This remains true when calculating the semiclassical contributions to the correlation function (25) as well since, to leading order in large k , the Wilson lines do not contribute to the saddle-point. Evaluating (25) in the background of the classical solution (18), the expectation value of Wilson lines in Chern-Simons gravity factorizes to leading order in large k

$$\langle W_{R_1}[\gamma_1] \cdots W_{R_n}[\gamma_n] \rangle_{CS} = \langle W_{R_1}[\gamma_1] \rangle_{CS} \cdots \langle W_{R_n}[\gamma_n] \rangle_{CS} + O\left(\frac{1}{k}\right) \quad (27)$$

Consequently, the factorization of the gravitational observables induces a factorization in the dual theory. Identifying the Chern-Simons coupling constant k with N^2 as in (23), (27) implies

$$\langle V_{R_1, \dots, R_n}[\gamma_1, \dots, \gamma_n] \rangle_{QFT} = \langle V_{R_1}[\gamma_1] \rangle_{QFT} \cdots \langle V_{R_n}[\gamma_n] \rangle_{QFT} + O\left(\frac{1}{N^2}\right) \quad (28)$$

which is the factorization property of gauge invariant observables in a large N rank gauge theory, as expected from very general considerations.

The purpose of these brief arguments is to motivate further investigation of the Chern-Simons gravity theory, and its supersymmetric extension [12, 13], for the purposes of gaining a better understanding of the relationship between generally covariant and quantum field theories. First and foremost, one would like to know more details about the conjectured dual quantum theory. This information may be obtained by a close investigation of explicit Wilson line calculations for the black hole background (18) we have considered, in analogy with calculations for $\mathcal{N} = 4$ super Yang-Mills theory [23]. It may even be possible to deduce a ‘loop equation’ for path-dependent objects in the dual theory from the Chern-Simons action.

Acknowledgements

We would like to thank M. Costa, M. Gutperle, M. Henningson, I. Klebanov, L. Smolin and D. Waldram for discussions. This work was completed with the support of the Natural Sciences and Engineering Research Council of Canada, NSF grant PHY98-02484 and the James S. McDonnell Foundation grant No. 91-48.

References

- [1] G. t'Hooft, “Dimensional Reduction in Quantum Gravity”; gr-qc/9310026.
- [2] L. Susskind, “The World as a Hologram” J. Math. Phys. **36** (1995), 6377; hep-th/9409089.
- [3] R. Bousso, “Holography in General Space-times” JHEP 9906 (1999) 028; hep-th/9906022.
- [4] I.R. Klebanov, “World Volume Approach to Absorption by Nondilatonic Branes” Nucl. Phys. **B496** (1997) 213; hep-th/9702076.
- [5] S.S. Gubser, I.R. Klebanov and A.A. Tseytlin, “String Theory and Classical Absorption by Three-branes” Nucl. Phys. **B499** (1997) 217; hep-th/9703040.
- [6] J. Maldacena, “The Large N Limit of Superconformal Field Theories and Supergravity” Adv. Theor. Math. Phys. **2** (1998) 231; hep-th/9711200.
- [7] S. Gubser, I. Klebanov and A. Polyakov, “Gauge Theory Correlators from Non-Critical String Theory” Phys. Lett. **B428** (1998) 105; hep-th/9802109.
- [8] E. Witten, “Anti-de Sitter Space and Holography” Adv. Theor. Math. Phys. **2** (1998) 253; hep-th/9802150.
- [9] O. Aharony, S.S. Gubser, J. Maldacena, H. Ooguri and Y. Oz, “Large N Field Theories, String Theory and Gravity”; hep-th/9905111.
- [10] L. Susskind and E. Witten, “The Holographic Bound in Anti-de Sitter Space”; hep-th/9805114.
- [11] A.H. Chamseddine, “Topological Gauge Theory of Gravity in Five and All Odd Dimensions” Phys. Lett. **B233** (1989) 291.
- [12] A.H. Chamseddine, “Topological Gravity and Supergravity in Various Dimensions” Nucl. Phys. **B346** (1990) 213.
- [13] M. Bañados, L.J. Garay and M. Henneaux, “The Dynamical Structure of Higher Dimensional Chern-Simons Theory” Nucl. Phys. **B476** (1996) 611; hep-th/9605159.

- [14] A. Achucarro and P.K. Townsend, “A Chern-Simons Action for Three-dimensional anti-de Sitter Supergravity Theories” *Phys.Lett.* **B180** (1986) 89.
- [15] E. Witten, “(2+1)-Dimensional Gravity as an Exactly Soluble System” *Nucl. Phys.* **B311** (1988) 46.
- [16] J. Gegenberg and G. Kunstatter, ‘Boundary Dynamics of Higher Dimensional AdS Space-time’; hep-th/9905228.
- [17] E. Witten, “Quantum Field Theory and the Jones Polynomial” *Comm. Math. Phys.* **121** (1989) 351.
- [18] S. Hawking and D. Page, “Thermodynamics of Black Holes in Anti-de Sitter Space” *Comm. Math. Phys.* **87** (1983) 577.
- [19] M. Bañados, C. Teitelboim and J. Zanelli, “Dimensionally Continued Black Holes” *Phys. Rev.* **D49** (1994) 975; gr-qc/9307033.
- [20] D.G. Boulware and S. Deser, “String-Generated Gravity Models” *Phys. Rev. Lett.* **55** (1985) 2656;
J.T. Wheeler, “Symmetric Solutions to the Gauss-Bonnet Extended Einstein Equations” *Nucl. Phys.* **B268** (1986) 737;
D.L. Wiltshire, “Spherically Symmetric Solutions of Einstein-Maxwell Theory with a Gauss-Bonnet Term” *Phys. Lett.* **169B** (1986) 36; “Black Holes in String-Generated Gravity Models” *Phys. Rev.* **D38** (1988) 2445.
- [21] R.C. Myers and J.Z. Simon, “Black-hole Thermodynamics in Lovelock Gravity” *Phys. Rev.* **D38** (1988) 2434; “Black Hole Evaporation And Higher Derivative Gravity” *Gen. Rel. Grav.* **21** (1989) 761;
B. Whitt, “Spherically Symmetric Solutions of General Second-Order Gravity” *Phys. Rev.* **D38** (1988) 3000;
T. Jacobson and R. C. Myers, “Entropy of Lovelock Black Holes” *Phys. Rev. Lett.* **70** (1993) 3684; hep-th/9305016.
- [22] G.W. Gibbons and S.W. Hawking, “Action Integrals and Partition Functions in Quantum Gravity” *Phys. Rev.* **D15** (1977) 2752.

- [23] E. Witten, “Anti-de Sitter Space, Thermal Phase Transition, and Confinement in Gauge Theories” *Adv. Theor. Math. Phys.* **2** (1998) 505; hep-th/9803131.
- [24] R. Troncoso and J. Zanelli, “New gauge supergravity in seven and eleven dimensions,” *Phys. Rev.* **D58**, 101703 (1998); hep-th/9710180.
- [25] P. Hořava, “M-Theory as a Holographic Field Theory” *Phys. Rev.* **D59** (1999) 46004; hep-th/9712130.
- [26] E. Witten, “Topology-Changing Amplitudes in 2+1 Dimensional Gravity” *Nucl. Phys.* **B223** (1989) 113.
- [27] S. Carlip, “Exact Quantum Scattering in 2+1 Dimensional Gravity” *Nucl. Phys.* **B324** (1989) 106.